

## Superior UPTAKE Zinc

Superior UPTAKE Zinc provides rapid foliar uptake of zinc to effectively correct and prevent deficiencies.

Superior UPTAKE Zinc is an advanced blend of organically chelated zinc, fulvates and amino acids. It is used to rapidly correct and prevent zinc deficiency in cereals, grain legumes, pasture legumes, oilseeds and fodder crops. Applied as a soil drench (via streaming nozzles) or as a foliar spray. Zinc is essential for the transformation of carbohydrates, and regulation and consumption of sugar in the plant.

## APPLICATION RATES

### Agricultural Crops and Pastures:

**Foliar:** Apply in early stages of growth when leaf area is sufficient to intercept spray. In high volume sprays, do not use more than a 1% solution; in low volume sprays (<50L/ha) use no more than 3% concentration. Repeat applications as required.

**Broad-acre Crops:** 1.0 - 2.0 litres/ha.

**Cereal Crops:** 0.5 - 1.5 litres/ha.

**Soil Applications:** Apply as required; do not exceed 10 litres/ha per application. Repeat applications as required.

## DIRECTIONS FOR USE

Superior UPTAKE Zinc is used in a wide range of agricultural crops and pastures.

Suitable for use as a soil drench (via streaming nozzles) and foliar spray. Applied through standard spraying equipment; always dilute prior to application.

# SUPERIOR UPTAKE ZINC

**Enhanced foliar uptake fertiliser**  
For broad-acre agriculture use

Rapidly correct and prevent zinc deficiency in broad-acre crops and pastures.

### Typical Analysis (wt/vol)

Zn- 15%, K- 0.2%, S- 7.4%, N- 0.1%,  
plus fulvates and amino acids.

## PRECAUTIONS

To prevent damage to crops:

- Spray during early morning or late afternoon; avoid hotter parts of the day or when crop is under heavy stress.
- Superior Uptake Trace Elements are physically compatible with a wide range of commonly used spray chemicals. Always mix a small quantity (eg. jar test) and check for residue in mixture. Do not continue if residue (sediment) occurs.
- Spray should not form visible droplet coalescence (DO NOT spray to run off).
- Spray under optimum crop growth conditions.

**NOTE:** If uncertain about spraying, we suggest a small area should be sprayed and observed prior to the total crop application (eg. when spraying for the first time or in combination with other chemicals).

**Wear protective clothing, safety glasses and gloves when using this product.**

## CLEAN UP PROCEDURE

Use all mixture in spray tanks; purge tanks and lines with clean water; flush lines. Do not return mix to original drums.

## STORAGE

Store in original container away from direct sunlight.

**FREE PLANT TISSUE REPORT**  
CALL 08 9384 3789

**Sustainable Ventures Pty Ltd**  
PO Box 419 Cottesloe WA  
Tel: (08) 9384 3789  
sustainablefarming@bigpond.com  
© Sustainable Ventures Pty Ltd 2009

Sustainable Ventures Pty Ltd has no control over farming methods, application of the product, weather or growing conditions. Consequently, no responsibility or liability for any failure in performance, losses, damages or injuries will be accepted by Sustainable Ventures Pty Ltd, its distributors or agents following the use of this product. Seek independent advice prior to use.

20L  200L  1000L