

## DIRECTIONS FOR USE

When mixed, fill spray tank half full with water and add the required amount of Organic-N while continuing to fill tank with water. Continue agitation through the spray operation. See Precautions. Always agitate drum contents before use.

## APPLICATION RATES

### Horticultural Crops:

Minimum dilution 1 : 100 with water.

(1: 150 for Almonds, Avocados, Citrus)

Foliar Spray: 2 - 5L/ha; repeat as required.

Irrigation: 5 - 10L/ha; repeat as required.

### Broad-acre Crops and Pastures:

Minimum dilution 1 : 30 with water.

Foliar Spray: 2-5L/ha; repeat as required.

Each farmer's situation and processes and practices may differ and therefore necessitate corrections to ensure optimum results. Adjust rates according to production levels. Repeat as required.

Always test for compatibility and crop suitability prior to use. Seek advice for rates suited to your crop.

## TYPICAL ANALYSIS

Nitrogen 10% w/v as amino acids.

With fulvic acid 29% w/v, trace elements (Ca, Mg, Zn, Fe), Organic Carbon 44% w/v.

## CLEAN UP PROCEDURE

Use all mixture in spray and irrigation tanks; purge tanks and lines with clean water; flush irrigation lines. Do not return mix to original drums.

FERTILISER - LIQUID

# Organic-N

Concentrated Liquid N - 10%

- For managing crop nitrogen demand and building yield
- Provides immediate nutrition for your crop
- Ideal to stimulate plant vegetative growth
- With 29% fulvic acid to improve nutrient uptake
- 100% soluble liquid
- Use via irrigation or foliar spray



□ 20L



□ 200L



□ 1,000L



## STORAGE

Store in original container away from direct sunlight. Store between 3°C and 40°C; a small amount of solid material may precipitate outside this temperature range. This does not affect efficacy of the product.

## PRECAUTIONS

Physically compatible with a wide range of commonly used products. Always mix a small quantity (jar test) and check for physical compatibility before combining with other ingredients.

When applying for the first time, or in combination with other products, a small test area should be sprayed and observed prior to the total spray.

Caution should be used when applying as a foliar spray in conjunction with copper or sulphur based products as increased activity of copper /sulphur may produce unacceptable levels of phytotoxicity.

## WARNING

Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with eyes and skin. Read SDS prior to use.

Wear protective clothing, safety glasses and gloves when using this product.

## DISCLAIMER

Sustainable Farming Solutions (SFS) has no control over farming methods, application of the product, irrigation systems, weather or growing conditions. Consequently no responsibility or liability for any failure in performance, losses, damages or injuries will be accepted by Sustainable Ventures Pty Ltd trading as SFS, its distributors or agents following the use of this product. Seek independent advice prior to use.



## Sustainable Farming Solutions

Tel: 08 9388 3623 : 03 9008 6352

admin@sustainablefarming.com.au

[www.sustainablefarming.com.au](http://www.sustainablefarming.com.au)