

## DIRECTIONS FOR USE

To mix, fill spray tank half full with water and add the required amount of Amino Power while continuing to fill tank with water. Continue agitation through the spray operation.

## APPLICATION RATES

### Horticultural Crops:

Foliar Spray: 250-500g/100L; repeat as required.

Irrigation: 2.5-5kg/ha; repeat as required.

### Broad-acre Crops and Pastures:

Foliar Spray: 250-500g/100L; repeat as required.

Each farmer's situation and processes and practices may differ and therefore necessitate corrections to ensure optimum results. Adjust rates according to production levels. Repeat as required.

Always test for compatibility and crop suitability prior to use. Seek advice for rates suited to your crop.

## TYPICAL ANALYSIS

Nitrogen 16% as amino acids.

## CLEAN UP PROCEDURE

Use all mixture in spray and irrigation tanks; purge tanks and lines with clean water; flush irrigation lines.

## STORAGE

Store in original container away from direct sunlight and moisture.



# FERTILISER - POWDER

# Amino Power

## Soluble Amino Acid Powder

- For managing crop nitrogen demand and building yield
- 16% Nitrogen
- 100% soluble fertiliser
- Use via irrigation or foliar spray

Manufacturing Date:

Batch #

25kg



## WARNING - NUISANCE DUST

Causes eye irritation.

May cause respiratory irritation.



Amino Acid component becomes sticky with moisture and can temporarily irritate eye.

Avoid breathing dust. Powder is fine and may cause irritation to respiratory tract.

Use a particle mask, safety glasses and protective clothing & gloves.

Wash skin thoroughly after handling. Avoid breathing dust.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if easy to do. Continue rinsing for 15 minutes; if eye irritation persists seek medical attention.

IF INHALED remove victim to fresh air. Call Poisons Information Centre 13 11 26 if you feel unwell.

See SDS for further first aid and safety instructions.

## PRECAUTIONS

Physically compatible with a wide range of commonly used products. Always mix a small quantity (jar test) and check for physical compatibility before combining with other ingredients.

When applying for the first time, or in combination with other products, a small test area should be sprayed and observed prior to the total spray.

Caution should be used when applying as a foliar spray in conjunction with copper or sulphur based products as increased activity of copper /sulphur may produce unacceptable levels of phytotoxicity.

## DISCLAIMER

Sustainable Farming Solutions (SFS) has no control over farming methods, application of the product, irrigation systems, weather or growing conditions. Consequently no responsibility or liability for any failure in performance, losses, damages or injuries will be accepted by Sustainable Ventures Pty Ltd trading as SFS, its distributors or agents following the use of this product. Seek independent advice prior to use.

## Sustainable Farming Solutions

Tel: 08 9388 3623 ; 03 9008 6352

admin@sustainablefarming.com.au

[www.sustainablefarming.com.au](http://www.sustainablefarming.com.au)